

SCREEN EXPOSURE AS A MODIFIABLE RISK

Pobitra Mishra , Medical Student , Samarkand state medical university, Uzbekistan

Misbahul Chaudhary , Medical Student , Samarkand state medical university,
Uzbekistan

Aravinthava , Medical Student , Samarkand state medical university, Uzbekistan

Hameed firnaz , Medical Student , Samarkand state medical university, Uzbekistan

Abstract

Digital screens have become part of everyday life, supporting education, communication, entertainment and access to health information. At the same time, prolonged or poorly timed screen exposure may displace sleep, physical activity, face-to-face interaction and other behaviours that protect health. This narrative review discusses screen exposure as a potentially modifiable risk factor, with particular attention to children and adolescents. Evidence is considered across sleep, physical activity and weight-related outcomes, vision and posture, and psychosocial well-being. The literature does not support treating every screen activity as harmful or assuming that screen duration alone explains health outcomes. Effects appear to depend on age, content, context, timing, the degree of adult involvement and what screen use replaces. Educational and socially supportive uses can be beneficial, while automatic playback, constant notifications and bedtime use can make balanced habits harder to maintain. Practical approaches include family media plans, screen-free meals and bedrooms, a pause from screens before sleep, regular movement breaks, co-viewing with younger children and age-appropriate content selection. The review concludes that screen exposure is best addressed through proportionate, family-centred and context-sensitive strategies rather than blame or a single rule for every child. Because this is a narrative review, the discussion is descriptive and does not constitute a systematic assessment or meta-analysis.

Keywords: screen exposure; screen time; children; adolescents; sleep; sedentary behaviour; digital health; prevention

1. Introduction

Phones, tablets, computers, televisions and gaming devices are now woven into ordinary routines. A child may use a tablet to practise reading, watch a cartoon while a caregiver prepares dinner, speak to relatives in another country and later play a game before bed. These activities look similar when measured as minutes in front of a screen, yet they differ in purpose, content, social setting and likely effect. For this reason, screen exposure should be understood as a pattern of behaviour rather than a single number.

The term “modifiable risk” means that exposure can be changed, at least partly, through personal choices, family routines, school practices and the design of digital environments. It does not mean that screens are inherently dangerous or that families can control every influence. Digital media can help children learn, create, maintain relationships and access support. The concern arises when use becomes excessive, difficult to stop, unsuitable for a child’s developmental stage or regularly replaces essential activities such as sleep, movement, meals and conversation.

Research reviews have reported associations between higher screen exposure and several health outcomes, including adiposity, depressive symptoms, diet quality and sleep. However, the strength of evidence varies by outcome, and many studies are observational. This means that an association does not prove that screen use alone caused the outcome. Family stress, neighbourhood safety, socioeconomic circumstances, pre-existing health difficulties and the content being viewed may influence both screen habits and health. A balanced review must therefore distinguish plausible mechanisms from established causal conclusions.




Figure 1. Conceptual illustration of shared, supervised screen use in a family setting.

2. Literature overview and approach

This narrative article draws on influential guidance and review literature identified through targeted searches of PubMed-indexed material, the American Academy of Pediatrics (AAP), the World Health Organization (WHO) and peer-reviewed review articles. Key sources include a systematic review of reviews on screen time and child and adolescent health, AAP guidance on media use and family media planning, and WHO guidance on sedentary behaviour. The aim is to explain recurring findings and practical implications, not to provide a reproducible systematic review.

An important limitation in this field is measurement. Some studies count television only; others include schoolwork, social media, video calls, gaming and background television. Self-reported time can be inaccurate, and the same duration may involve very different levels of attention or physical inactivity. Evidence also changes as platforms and devices evolve. Findings should therefore be interpreted as a broad picture rather than as a precise prediction for an individual child.

3. Potential pathways linking screen exposure and health

3.1 Sleep and daily rhythm

Screen use can affect sleep in several ways. A video or game may delay bedtime because the user wants to continue; notifications can interrupt wind-down routines; and stimulating content can make it harder to settle. Light from a screen may also contribute to circadian effects, although real-world sleep outcomes depend on brightness, timing, duration and individual sensitivity. Sleep matters for attention, mood, learning, growth and general well-being, so a small nightly delay repeated across a week may become meaningful.

Practical advice should focus less on fear of a device and more on protecting a predictable bedtime. Families can charge phones outside bedrooms, disable non-essential notifications and agree on a screen-free period before sleep. AAP guidance commonly recommends avoiding screens during the hour before bedtime and keeping devices out of sleeping spaces when possible. These steps are easier to follow when adults adopt them too.

3.2 Sedentary behaviour, diet and weight

Many screen activities involve sitting for long periods. When they replace outdoor play, walking or sport, daily movement can fall. Television and online content may also encourage distracted eating or expose children to advertising for energy-dense foods. These pathways can help explain why research has found associations between greater screen exposure and higher adiposity or less healthy dietary patterns. Still, screen use is not the only driver of weight: food access, sleep, genetics, medication, family routines and the built environment also matter.

A useful goal is to interrupt long sitting periods rather than simply counting minutes. A short walk, stretching, dancing to music or active play between online tasks can make a difference to the daily routine. Meals and snacks are good opportunities to put devices away and notice hunger and fullness cues.

3.3 Vision, headaches and posture

Extended near work can contribute to eye strain, dry or tired eyes, blurred vision and headaches, particularly when a person blinks less, holds a device very close or uses it continuously without breaks. Prolonged downward head posture and poorly arranged desks may also contribute to neck, shoulder or back discomfort. These symptoms are not unique to digital screens, and persistent pain or visual changes should not automatically be attributed to screen exposure.

Simple adjustments include using comfortable lighting, placing the screen at a reasonable distance, increasing text size, taking regular visual breaks and changing posture. The commonly used 20-20-20 reminder—every 20 minutes, look at something about 20 feet away for 20 seconds—is a practical habit rather than a guarantee against eye problems. Outdoor time is also valuable, particularly for children, and should not be replaced by additional screen-based leisure.

3.4 Emotional and social well-being

Digital media can support friendship, identity, creativity and connection, especially for young people who feel isolated offline. It can also expose users to cyberbullying, unrealistic comparisons, distressing content, privacy risks or pressure to remain constantly available. The impact is shaped by what a person sees, how they feel during and after use, and whether online activity supports or disrupts their relationships and responsibilities.

Parents and educators should look for changes in functioning rather than assume that every frequent user has a problem. Warning signs may include persistent sleep loss, falling school participation, abandoning previously enjoyed activities, repeated conflict about stopping, or distress linked to online interactions. Calm conversations and practical support are generally more helpful than shame or sudden punishment.

4. Summary of evidence and practical interpretation

Area	What research suggests	Practical interpretation
Sleep	Higher or bedtime screen use is associated with poorer sleep outcomes in some studies.	Protect bedtime routines; keep devices away from beds when feasible.
Weight and diet	Review-level evidence links greater screen time with adiposity and less healthy diet patterns.	Pair device boundaries with movement, regular meals and healthy food access.
Mental well-being	Associations with depressive symptoms are reported; findings vary by type of use and person.	Ask about content, online experiences and functioning, not time alone.
Vision and posture	Continuous near work may cause temporary strain or discomfort.	Use breaks, comfortable positioning and appropriate lighting; assess persistent symptoms.
Learning and connection	High-quality media can support learning and relationships; passive or distracting use may interfere with tasks.	Choose age-appropriate content and use media together when useful.

Table 1. Broad evidence themes and practical implications. These statements summarise associations and guidance; they should not be read as proof that screen exposure alone causes each outcome.

5. Screen exposure as a modifiable risk: what can be changed?

Effective prevention is usually realistic, consistent and shared. A strict rule that ignores school requirements, family work schedules or a child's needs may fail quickly. Instead, families can identify the situations where screens are most likely to displace something important, then make one or two changes at a time. For example, a household may start by keeping meals device-free and charging phones outside bedrooms, then review whether sleep and family interaction improve.




Figure 2. Four practical pillars for healthier screen habits: sleep, movement, connection and content choice.

5.1 Guidance by developmental stage

For very young children, real-world interaction, play, language-rich conversation and sleep are especially important. AAP guidance advises avoiding screen media before 18 months except video chatting, choosing high-quality content and watching together for children aged 18–24 months, and limiting use to about one hour per day of high-quality programming for children aged 2–5 years. Recommendations can differ slightly across organisations and are updated as evidence develops; caregivers should consult current local guidance and their child's clinician when needed.

For school-age children and adolescents, a single daily limit is often less useful than a plan that protects sleep, physical activity, school responsibilities and in-person relationships. Families can discuss when screens are useful, which activities are recreational, what content is appropriate and what to do when online experiences become upsetting. The plan should be reviewed as a child's maturity, school demands and technology use change.

5.2 A sample family media plan

Routine	Example agreement
Meals	Phones and tablets stay away from the table.
Bedtime	Devices charge outside bedrooms; quiet wind-down begins about one hour before sleep.
Study	School tasks are done with non-essential notifications off.
Movement	Stand, stretch or move between long sitting periods.
Young children	Adults choose suitable content and watch or play together where possible.
Review	Family checks the plan weekly and changes rules that are unrealistic.

Table 2. Illustrative family agreement. It should be adapted to age, accessibility needs, school demands and family circumstances.

5.3 Schools, clinicians and digital platforms

Schools can help by teaching digital literacy, building movement into the day and avoiding unnecessary device use when a non-digital method serves the same purpose. Clinicians can ask non-judgmental questions about sleep, mood, physical activity and media habits during routine visits. Platforms and device makers also have a role: clear parental controls, age-appropriate defaults, reduced manipulative notifications and tools that make stopping easier can support healthier behaviour. Prevention should not place the entire responsibility on individual parents when digital products are designed to hold attention.

6. Discussion

The strongest public-health message is not that all screen exposure is harmful, but that its timing, content and opportunity cost matter. A child video-calling a grandparent, completing a guided science lesson or creating digital art is not having the same experience as a child watching autoplay videos late into the night. Measuring total minutes alone can hide these differences. A more useful assessment asks what the screen is being used for, whether the child can stop, what activity it replaces and how the child feels afterwards.

Evidence also needs careful interpretation. Much of the literature is observational, so direction of effect can be complicated. For example, poor sleep may lead to more nighttime device use, as well as nighttime device use contributing to poor sleep. Similarly, a young person experiencing low mood may seek online distraction or support. Longitudinal studies, objective device-use measures and studies that distinguish content and context can improve understanding. Future research should include diverse family settings and children with disabilities, for whom digital access may provide essential communication or participation.

Screen exposure is nevertheless a practical prevention target because routines can be adjusted without eliminating technology. Small changes—such as moving a charger out of the bedroom, turning off autoplay, taking movement breaks and making meals screen-free—are low-cost and can be tested at home. The best plan is one that improves daily life and can be maintained, rather than one that creates constant conflict.

7. Conclusion

Screen exposure is a modifiable influence on health, but its effects are neither uniform nor determined by time alone. Higher or poorly timed use may contribute to sleep disruption, sedentary routines, less healthy eating, temporary eye strain and some psychosocial difficulties, while purposeful digital media can support learning and social connection. Families, schools, clinicians and technology providers can reduce avoidable risks by protecting sleep, encouraging movement, prioritising face-to-face relationships, choosing suitable content and revisiting clear media boundaries. A balanced, developmentally appropriate approach is more defensible than fear-based messaging or a universal rule for every child.

References

1. Stiglic N, Viner RM. Effects of screentime on the health and well-being of children and adolescents: a systematic review of reviews. *BMJ Open*. 2019;9(1):e023191. doi:10.1136/bmjopen-2018-023191.
2. American Academy of Pediatrics, Council on Communications and Media. Media Use in School-Aged Children and Adolescents. *Pediatrics*. 2016;138(5):e20162592. doi:10.1542/peds.2016-2592.
3. American Academy of Pediatrics. The Role of the Pediatrician in the Promotion of Healthy, Active Living. *Pediatrics*. 2024;153(3):e2023065480. Available from: <https://publications.aap.org/pediatrics/article/153/3/e2023065480/>

4. World Health Organization. WHO Guidelines on Physical Activity and Sedentary Behaviour. Geneva: World Health Organization; 2020. Available from: <https://www.who.int/publications/i/item/9789240015128>
5. American Academy of Pediatrics. How to Make a Family Media Plan. HealthyChildren.org. Updated January 20, 2026. Available from: <https://www.healthychildren.org/English/family-life/Media/Pages/How-to-Make-a-Family-Media-Use-Plan.aspx>
6. Moreno MA, Radesky J, Walsh MC, Tomopoulos S. The Family Media Plan. Pediatrics. 2024;154(6):e2024067417. doi:10.1542/peds.2024-067417.
7. American Academy of Pediatrics. Helping Kids Thrive in a Digital World: AAP Policy Explained. HealthyChildren.org. Updated June 3, 2026. Available from: <https://www.healthychildren.org/English/family-life/Media/Pages/helping-kids-thrive-in-a-digital-world-AAP-policy-explained.aspx>
8. Digital screen exposure in infants, children and adolescents: a systematic review of existing recommendations. PubMed record. 2025. Available from: <https://pubmed.ncbi.nlm.nih.gov/41127851/>