

GUARDING THE ENGINE OF LIFE**Modifiable and Non-Modifiable Causes of Cardiac Problems — and How Prevention Changes the Story**

Majid Reza , Medical Student, Samarkand state medical university, Uzbekistan

Asif Eqbal , Medical Student, Samarkand state medical university, Uzbekistan

Kumar Aakash , Medical Student, Samarkand state medical university, Uzbekistan

Hussain Rokibul , Medical Student, Samarkand state medical university, Uzbekistan

Abstract

Cardiovascular disease remains the single leading cause of death worldwide, claiming close to 18 million lives each year, yet a large share of this burden is neither inevitable nor mysterious. This narrative article reviews the causes of cardiac problems through a practical lens, sorting them into non-modifiable factors — age, sex, family history and heredity, and ethnic background — which cannot be changed but can be anticipated, and modifiable factors — tobacco use, high blood pressure, abnormal cholesterol, diabetes and obesity, physical inactivity, unhealthy diet, harmful alcohol use, chronic stress, and poor sleep — which respond to everyday choices and medical care. Drawing on landmark studies such as the Framingham Heart Study and the INTERHEART study, the article argues that roughly nine in ten heart attacks are explained by modifiable risk factors acting together, which means prevention is not a matter of luck but of strategy. Tables summarize the distinction between the two categories and the clinical targets for each modifiable factor, while figures illustrate the anatomy at stake and the lifestyle levers that protect it. The central message is hopeful: we cannot choose our genes or our birth year, but we can decide, to a remarkable degree, how our hearts will age.

Keywords: cardiovascular disease; risk factors; modifiable; non-modifiable; prevention; lifestyle; hypertension; atherosclerosis

1. Introduction

Every day, the human heart beats around 100,000 times, quietly pumping nearly 7,500 litres of blood through a network of vessels that would stretch more than 100,000 kilometres if laid end to end. It asks for remarkably little in return — no conscious

effort, no special attention — which is perhaps why so many of us take it for granted until the day it falters.

The good news — and it is genuinely good news — is that a large part of that story is written by factors we can influence. Researchers divide the causes of heart disease into two broad families. The first, the non-modifiable causes, are the cards we are dealt: our age, our sex, the genes we inherit, and the ethnic community into which we are born. We cannot change these, but we can measure their weight and plan around them. The second family, the modifiable causes, includes behaviours and conditions such as smoking, blood pressure, cholesterol, blood sugar, body weight, diet, exercise, alcohol, stress, and sleep. These are the levers within our reach — and the evidence shows that pulling them, even modestly and even later in life, dramatically changes the odds.

2. The Heart Under Threat: A Brief Look at What Can Go Wrong

Most cardiac problems share a common villain: atherosclerosis, the gradual build-up of fatty plaques inside the walls of arteries. Imagine a water pipe that slowly collects rust and scale on its inner surface. For years the water still flows, and everything seems fine. Then one day a patch of scale cracks, a clot forms on top, and the pipe suddenly blocks. When the 'pipe' is a coronary artery feeding heart muscle, the result is a myocardial infarction — a heart attack. When it is an artery in the brain, the result is a stroke. Chronic high pressure in the system leads to heart failure; erratic electrical signals produce arrhythmias such as atrial fibrillation. Nearly every risk factor discussed below either accelerates plaque build-up, raises blood pressure against vessel walls, or disturbs the heart's rhythm and metabolism.




Figure 1. Surface anatomy of the human heart. The coronary arteries visible on its surface supply the muscle itself — they are the 'pipes' most often blocked in a heart attack. (Educational illustration.)

3. Non-Modifiable Causes: The Hand We Are Dealt

Non-modifiable risk factors cannot be removed by diet, exercise, or medication. Calling them 'causes' requires a small caution: with the partial exception of inherited disorders, they are better understood as amplifiers. They do not create plaque by themselves so much as they raise the baseline pressure of risk against which every other factor acts.

Age is the most powerful of all. The lifetime of a heart is written in its vessels, and decades of pulsatile blood flow leave their mark. The risk of a first heart attack roughly doubles with each decade after age 40, and most events occur after the age of 65. This is not a verdict of doom — it is a calendar. Knowing that risk climbs with age tells us that prevention should begin early, when habits are still forming and vessels are still healthy.

Sex shapes the risk profile in ways that biology and medicine are still untangling. Men generally develop coronary disease earlier than women; a man's lifetime risk is appreciably higher in middle age. Women are partly protected by oestrogen before menopause, yet their risk catches up and, after 65, can exceed men's. Worse, women are more likely to have atypical symptoms and to be underdiagnosed, which means their non-modifiable disadvantage is often compounded by a modifiable one — delayed care.

Family history is the factor that surprises people most. Having a first-degree relative — a parent or sibling — who suffered a heart attack or stroke before age 55 (men) or 65 (women) roughly doubles a person's own risk. At the extreme end of inheritance sit monogenic disorders such as familial hypercholesterolaemia, in which a single faulty gene keeps cholesterol dangerously high from childhood. None of this can be edited away, but it can be screened for, monitored, and aggressively managed — a family history, in effect, is an early-warning system, not a death sentence.

Ethnicity and geography complete the picture. South Asian communities, for instance, carry a markedly higher burden of coronary disease at younger ages and lower body weights, while some African and Afro-Caribbean populations show higher rates of hypertension and stroke. These differences reflect a tangled mix of genetics, diet, social conditions, and access to care — a reminder that 'non-modifiable' and 'unimportant' are very different things.

Non-modifiable factor	Why it raises cardiac risk	What CAN still be done
Age	Arterial wear accumulates over decades; risk roughly doubles each decade after 40.	Start screening early; manage every modifiable factor more strictly with age.
Sex	Men at higher risk earlier; women catch up after menopause and are often underdiagnosed.	Sex-aware symptom awareness; regular check-ups especially around menopause.
Family history / genetics	Inherited lipid disorders and shared genes can double risk; familial hypercholesterolaemia is a prime example.	Screen relatives; earlier and more intensive cholesterol and blood pressure control.
Ethnic background	Some groups carry higher susceptibility at younger ages or to specific conditions (e.g., hypertension).	Community-targeted screening; culturally adapted diet and lifestyle programmes.

Table 1. Non-modifiable causes of cardiac disease: they cannot be changed, but their consequences can be anticipated and blunted.

4. Modifiable Causes: Where Prevention Actually Lives

If the previous section is about fate, this one is about freedom. The modifiable causes of cardiac disease are the everyday variables — some behavioural, some clinical — that individuals, clinicians, and communities can genuinely alter. The INTERHEART study, which compared roughly 15,000 heart attack patients with an equal number of matched controls across 52 countries, produced the statistic that anchors this entire field: nine potentially modifiable risk factors together explained more than 90% of the population-attributable risk of a first heart attack in men, and around 94% in women. In plain language, the vast majority of heart attacks are, in principle, preventable.




Figure 2. The approximate share of first-heart-attack risk attributable to modifiable versus non-modifiable factors, based on the INTERHEART case-control study (Yusuf *et al.*, *Lancet*, 2004). Values are approximate population-attributable estimates.

Tobacco sits near the top of any honest list. Smoking damages the endothelium — the delicate inner lining of blood vessels — within minutes of the first puff, promotes clotting, lowers 'good' HDL cholesterol, and accelerates plaque growth. The harm is dose-related but also fast-reversible: within a year of quitting, heart disease risk falls by roughly half, and within fifteen years it approaches that of a never-smoker. Few actions in all of medicine repay themselves so quickly.

High blood pressure, the 'silent killer,' rarely announces itself until damage is done. Persistent pressure hardens and scars arteries, thickens the heart muscle, and sets the stage for both heart failure and stroke. Because it produces no symptoms for most people, it is discovered only by measurement — which is why the simple act of having one's blood pressure checked regularly is itself a preventive intervention.

Abnormal cholesterol — especially high LDL ('bad') cholesterol and triglycerides, and low HDL — supplies the raw material for atherosclerotic plaque. Modern guidelines no longer treat cholesterol as a stand-alone number but as one ingredient in an overall risk score, since the same LDL level means very different things for a 30-year-old woman and a 60-year-old man with diabetes.

Diabetes and obesity form a closely linked pair. Persistently high blood sugar injures small vessels and nerves, accelerates coronary disease, and roughly doubles cardiovascular risk; diabetes is now considered a 'coronary risk equivalent.' Obesity — particularly fat carried around the waist — feeds the problem through insulin resistance, inflammation, and strain on the heart. The encouraging corollary is that losing 5–10% of body weight produces measurable improvements in blood pressure, sugar, and lipids within months.

Physical inactivity, an unhealthy diet, harmful alcohol use, chronic stress, and poor sleep round out the list. Sedentary lifestyles weaken the heart's efficiency and worsen every metabolic number; diets heavy in salt, trans fats, processed meat, and sugar do the same from the inside. Alcohol's relationship with the heart is a J-shaped curve often misquoted: while very light drinking was once thought protective, recent large analyses find that the safest level for the cardiovascular system is none at all. Chronic psychological stress and depression keep the body flooded with stress hormones that raise pressure and promote inflammation, and short or fragmented sleep compounds every one of these effects — the neglected hours of the night turn out to be working hours for heart health.

Modifiable factor	Why it harms the heart	Evidence-based target
Smoking / tobacco	Endothelial damage, clotting, low HDL; risk ↑ even from a few cigarettes/day.	Complete cessation; risk falls ~50% within 1 year of quitting.
High blood pressure	Hardens arteries, thickens heart muscle → heart failure, stroke.	Generally < 130/80 mmHg in most adults (individualized by clinician).
High LDL cholesterol	Supplies the lipid core of atherosclerotic plaque.	LDL lowered per overall risk score; very high-risk patients aim < 55–70 mg/dL.
Diabetes / high blood sugar	Damages small vessels and nerves; ~doubles CVD risk.	HbA1c usually < 7% (individualized); tight BP + statin therapy for protection.
Obesity (esp. abdominal)	Insulin resistance, inflammation, cardiac workload.	BMI 18.5–24.9; waist < 94 cm (men) / < 80 cm (women); lose 5–10% weight for benefit.
Physical inactivity	Weak cardiac fitness; worsens BP, sugar, lipids, weight.	≥150 min/week moderate activity (e.g., brisk

		walking) or 75 min vigorous.
Unhealthy diet	Excess salt, trans fat, processed meat, sugar → BP, lipids, weight.	Vegetables, fruit, whole grains, legumes, fish; nuts in moderation; limit salt < 5 g/day.
Harmful alcohol use	Raises BP, arrhythmia risk, cardiomyopathy; no safe level for CVD.	Avoid or strictly limit; do not start 'for the heart.'
Chronic stress / depression	Stress hormones → hypertension, inflammation, poor self-care.	Regular relaxation, social connection, professional help when needed.
Poor sleep (< 7 h / apnea)	Linked to hypertension, obesity, arrhythmia, diabetes.	7–9 hours nightly; screen for and treat obstructive sleep apnoea.

Table 2. The modifiable causes at a glance: the damage mechanism and the practical target for each.

5. Prevention in Practice: Turning Knowledge into Habit

Knowing the causes is one thing; living the prevention is another. The clinical guidelines of the American Heart Association and the American College of Cardiology frame primary prevention around a simple hierarchy: first, assess total risk; second, treat the obvious villains (smoking, blood pressure, cholesterol, diabetes); third, build the lifestyle scaffold — diet, activity, weight, sleep — on which everything else rests. Risk calculators such as the pooled cohort equations convert a person's age, sex, pressures, and numbers into a ten-year probability, helping patient and doctor decide together how aggressive treatment should be.



Figure 3. Evidence-based pillars of cardiovascular prevention: nutrition, exercise, tobacco avoidance, social determinants, and informed medication use. (Infographic by the American Heart Association, 2019.)

6. Discussion: Reading Both Sides of the Balance Sheet Together

It would be a mistake to treat this article's two sections as separate lists. The non-modifiable and modifiable factors interact constantly, and their interaction is precisely where prevention earns its keep. A 58-year-old man with a strong family history (non-modifiable) who quits smoking, walks daily, and brings his blood pressure under control (modifiable) can easily carry a lower absolute risk than a 30-year-old with clean genes who smokes, sleeps five hours, and never checks a single number. The dealt cards matter — but the way the hand is played matters more.

This is also why 'non-modifiable' must never become an excuse for fatalism. Age cannot be reversed, but its vascular consequences can be slowed; a parent early heart attack cannot be undone, but it can trigger family screening that saves the next generation; ethnicity cannot be chosen, but culturally tailored community programmes demonstrably improve outcomes in high-risk groups. The honest reading of the evidence is not 'your fate is fixed' but 'your risk is readable' — and what can be read can, to a remarkable degree, be managed.

7. Review of the Literature

The scientific foundation for this article rests on several pillars of cardiovascular epidemiology. The Framingham Heart Study, launched in 1948 and following a Massachusetts town across generations, was the first large cohort to identify the major risk factors — cholesterol, blood pressure, smoking — as measurable, trackable predictors of coronary events, and to quantify how they compound one another. Framingham's enduring gift was the idea of 'risk factors' itself: the notion that a disease of middle age is forecast by measurable conditions of early adulthood.

Half a century later, the INTERHEART study brought this framework to the whole world. By enrolling cases and controls from every inhabited continent, it demonstrated that the same handful of modifiable factors — lipids, smoking, hypertension, diabetes, obesity, inactivity, diet, alcohol, and psychosocial stress — dominates heart attack risk across radically different populations, and that their combined population-attributable risk exceeds 90%. Global Burden of Disease analyses have since confirmed the picture at the level of entire nations, ranking high blood pressure, tobacco, unhealthy diet, and high blood sugar among the leading metabolic and behavioural drivers of premature death worldwide.

8. Conclusion

The causes of cardiac problems fall into two families, and honesty requires naming both. The non-modifiable causes — age, sex, heredity, and ethnicity — set the stage on which the drama is played, and they deserve respect, screening, and early attention. But the stage is not the script. The modifiable causes, which explain the overwhelming majority of preventable heart disease, are written daily: in what we smoke or refuse to smoke, what we eat, how we move, how we sleep, how we manage pressure — both the blood-borne and the life-borne kind. Prevention, in the end, is not a pill or a single dramatic resolution. It is the quiet, repeated arithmetic of small choices, made over years, adding up to a heart that keeps its rhythm for a lifetime. We cannot choose our genes. We can, with knowledge and persistence, choose a great deal of everything else.

References

1. World Health Organization. Cardiovascular diseases (CVDs) — Fact sheet. Geneva: WHO; 2021. Available from: [https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-\(cvds\)](https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-(cvds)).
2. Yusuf S, Hawken S, Ôunpuu S, Dans T, Avezum A, Lanas F, et al. Effect of potentially modifiable risk factors associated with myocardial infarction in 52

countries (the INTERHEART study): case-control study. *Lancet*. 2004;364(9438):937-52.

3. GBD 2019 Risk Factors Collaborators. Global burden of 87 risk factors in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet*. 2020;396(10258):1223-49.

4. Arnett DK, Blumenthal RS, Albert MA, Buroker AB, Goldberger ZD, Hahn EJ, et al. 2019 ACC/AHA Guideline on the Primary Prevention of Cardiovascular Disease. *Circulation*. 2019;140(11):e596-e646.

5. Kannel WB, Dawber TR, Kagan A, Revotskie N, Stokes J 3rd. Factors of risk in the development of coronary heart disease — six year follow-up experience: the Framingham Study. *Ann Intern Med*. 1961;55(1):33-50.

6. Ettehad D, Emdin CA, Kiran A, Anderson SG, Callender T, Emberson J, et al. Blood pressure lowering for prevention of cardiovascular disease and death: a systematic review and meta-analysis. *Lancet*. 2016;387(10022):957-67.

7. Ekelund U, Tarp J, Fagerland MW, Johannessen JS, Hansen BH, Jefferis BJ, et al. Joint associations of accelerometer-measured physical activity and sedentary time with all-cause mortality: a harmonised meta-analysis. *BMJ*. 2019;366:14570.

8. Wood AM, Kaptoge S, Butterworth AS, Willeit P, Warnakula S, Bolton T, et al. Risk thresholds for alcohol consumption: combined analysis of individual-participant data for 599,912 current drinkers in 83 prospective studies. *Lancet*. 2018;391(10129):1513-23.

9. Micha R, Peñalvo JL, Cudhea F, Imamura F, Rehm CD, Mozaffarian D. Association between dietary factors and mortality from heart disease, stroke, and type 2 diabetes in the United States. *JAMA*. 2017;317(9):912-24.

10. Lloyd-Jones DM, Braun LT, Ndumele CE, Smith SC Jr, Sperling LS, Virani SS, et al. Use of risk assessment tools to guide decision-making in the primary prevention of atherosclerotic cardiovascular disease: a special report from the American Heart Association and American College of Cardiology. *Circulation*. 2019;139(25):e1162-e77.

11. Peters SAE, Woodward M. Sex differences in the burden and complications of cardiovascular disease. *Nat Rev Cardiol*. 2015;12(4):231-41.

12. Khera AV, Chaffin M, Aragam KG, Haas ME, Roselli C, Choi SH, et al. Genome-wide polygenic scores for common diseases identify individuals with risk equivalent to monogenic mutations. Nat Genet. 2018;50(9):1219-24.