

OCCUPATIONAL HEALTH HAZARDS AMONG MEDICAL STUDENTS DURING CLINICAL TRAINING

Mohd Sohel SULEMANI , Medical Student, Samarkand state medical university,
Uzbekistan

Mohd Umar saifi , Medical Student, Samarkand state medical university, Uzbekistan

Mohammad bishr shakil , Medical Student, Samarkand state medical university,
Uzbekistan

Shanu saifi , Medical Student, Samarkand state medical university, Uzbekistan

Abstract

Medical students occupy an unusual position in the hospital: they are learners, yet they work in the same wards, theatres, and emergency rooms as fully trained staff, often with the least experience, the least authority, and the fewest protective habits. This narrative article reviews the occupational health hazards they face during clinical training and argues that these hazards are neither rare nor incidental, but structural. The hazards are organized into five families — biological (needlestick injuries and blood-borne pathogens such as HIV, hepatitis B and C, and tuberculosis), chemical (antineoplastic drugs, anaesthetic gases, disinfectants), physical (radiation, noise, slips and falls), ergonomic (patient handling, long standing hours), and psychological (sleep deprivation, burnout, depression, and bullying) — each illustrated with examples and, where possible, figures from the published literature. Tables summarize the hazard categories and the reasons students are disproportionately exposed. The article concludes that protecting students requires more than personal protective equipment: it demands vaccination, safety-engineered devices, enforced reporting systems, duty-hour limits, and above all a clinical culture in which asking for help is a sign of professionalism rather than weakness.

Keywords: medical students; clinical training; occupational health; needlestick injury; burnout; blood-borne pathogens; workplace safety

1. Introduction

There is a peculiar moment in the life of every medical student: the first time a white coat is handed over in a hospital corridor, and the realization dawns that the coat is less a uniform than a passport — into other people's illnesses, other people's emergencies, and, quietly, other people's risks. From that day forward, the student is exposed to the same biological, chemical, physical, and emotional hazards as the professionals beside them, but without their years of conditioning, their instincts, or sometimes even their gloves.

Occupational health is the discipline concerned with preventing work-related injury and illness, and for two centuries it has asked the same sensible question of every workplace: what does this environment do to the people inside it? Hospitals, oddly, have been slow to turn that question inward. We screen patients meticulously for infections we might transmit; we audit hand hygiene with near-religious devotion; yet the tally of injuries suffered by the students themselves — the needlestick in a crowded casualty bay, the needle prick during a nervous first venepuncture, the month of night shifts that erases sleep and mood alike — has historically been kept less carefully than the ward inventory.

This article examines the occupational health hazards of medical students during clinical training in a narrative rather than purely technical style. It walks through the major categories of hazard, pausing on the ones the literature identifies as most common and most consequential, and then asks why students are so consistently overexposed. It closes with prevention — because every hazard described here already has a countermeasure, and the real scandal is how rarely those countermeasures are applied with consistency.

2. The Clinical Years: When the Classroom Meets the Ward

The pre-clinical years carry their own pressures, but it is the clinical phase — ward rotations, operating theatres, emergency departments, and community postings — that converts theoretical risk into daily exposure. A student drawing blood for the first time performs a procedure that staff perform hundreds of times a month; a student standing through a twelve-hour theatre list absorbs the same anaesthetic gases, radiation scatter, and fatigue as the surgical team, but with no say in the schedule. The teaching hospital is, in effect, a live laboratory in which students are both observers and, increasingly, participants. That participation is precisely what makes training valuable — and what obliges institutions to treat student safety as a duty rather than an afterthought.



*Figure 1. Medical students in the clinical environment — wards and theatres are where classroom learning becomes hands-on exposure to occupational hazards.
(Photo: in-Training, online publication for medical students.)*

3. Biological Hazards: Needles, Blood, and Invisible Enemies

Ask any group of senior clinicians about their training and a surprising number will roll up a sleeve or hold out a hand to show a small pale scar — the signature of a needlestick injury sustained as a student. The global numbers justify the storytelling: it has been estimated that sharps injuries expose millions of health workers to blood-borne pathogens every year, and large cohort studies have found that trainees sustain a disproportionate share. The mechanics are unglamorous — a needle passed from nurse to doctor between patient curtains, a syringe recapped in a hurry, a sharps box that overflows because nobody checked it — but the consequences can be life-altering. A single hollow-bore needle from an infectious patient can transmit hepatitis B in up to roughly a third of susceptible exposures, hepatitis C in a smaller but still substantial fraction, and HIV in around one in three hundred. Those odds sound comforting until one remembers that students perform and assist at hundreds of procedures, often on patients whose infection status is unknown.

Beyond sharps, students face the same airborne and contact risks as any hospital worker: tuberculosis from the undiagnosed cough in a crowded outpatient department, influenza and — as the COVID-19 pandemic made brutally clear — novel respiratory pathogens for which the student may lack adequate personal protective equipment. Latex allergy, scabies, and drug-resistant organisms complete the biological portfolio. The recurring theme is unpredictability: the student rarely knows which patient carries which risk, and is often the last person in the chain to be told.

4. Beyond Infection: Chemical, Physical, Ergonomic, and Psychological Hazards

The word 'hazard' in a hospital summons images of needles, but some of the most stubborn occupational threats carry no syringe. Antineoplastic drugs handled during oncology rotations can be absorbed through skin or inhaled as dust; glutaraldehyde and other disinfectants irritate airways; waste anaesthetic gases accumulate in poorly ventilated theatres. Physically, students absorb scattered radiation during fluoroscopy lists, endure noise levels that would trouble an industrial inspector, and navigate slippery corridors on too little sleep. Ergonomically, the profession seems almost designed to injure the young: hours of standing, awkward positioning at operating tables, and patient transfers performed with technique learned by imitation rather than instruction.

The psychological hazards, however, may be the heaviest of all. Systematic reviews report that roughly a quarter to a third of medical students screen positive for depressive symptoms, while burnout rates climb higher still as training progresses. The drivers are structural: chronic sleep deprivation on call nights, the fear of making a fatal mistake, examinations that never quite end, and — too often — humiliation from seniors who mistake cruelty for teaching. Add exposure to death, suffering, and occasional abuse from patients, and it becomes clear that the clinical years tax the mind as reliably as the body.




Figure 2. Approximate prevalence ranges for selected hazards among medical students, as reported across published studies; figures vary widely by country and measurement instrument and are illustrative rather than definitive.

Hazard category	Representative examples	Potential consequences
-----------------	-------------------------	------------------------

Biological	Needlestick/sharps injuries; blood & body-fluid exposure; tuberculosis; influenza and other respiratory infections; latex	HIV, hepatitis B/C; chronic infection; TB disease; allergy
Chemical	Antineoplastic drugs; waste anaesthetic gases; glutaraldehyde and disinfectants; formalin	Organ toxicity; mucosal irritation; dermatitis; reproductive harm
Physical	Ionizing radiation; noise; wet floors; needles in laundry; electric equipment	Cumulative radiation dose; hearing strain; falls and lacerations
Ergonomic	Prolonged standing; awkward surgical postures; manual patient handling; heavy file/equipment loads	Musculoskeletal pain; varicose veins; chronic back injury
Psychosocial	Sleep deprivation; examination pressure; bullying; witnessing death; work–study imbalance	Depression, anxiety, burnout; reduced empathy; impaired learning

Table 1. The five families of occupational hazards encountered by medical students during clinical training.

5. Why Students Are Disproportionately Exposed

It would be tempting to dismiss student injuries as the ordinary tuition of a dangerous craft. The evidence suggests the opposite: students are exposed not merely because they work in risky places, but because of who they are within those places. They are inexperienced, performing procedures for the first time under observation, often with unfamiliar equipment and in settings where supervision competes with service demands. They occupy the bottom of an unforgiving hierarchy: the student who asks for gloves, questions a sharps practice, or refuses an unsafe task risks being labelled difficult at precisely the moment they need referees for their careers. Reporting systems, where they exist, are frequently designed for employees rather than students, so incidents go unrecorded, unprophylaxed, and uncounted. And because students rotate between departments and hospitals, nobody's dashboard accumulates their total exposure. Each of these factors is fixable; together, they explain why the newest hands in the hospital so often collect its oldest hazards.

Key exposure	Why students are at higher risk	Core protective measure
Needlestick / sharps injury	First-time, supervised procedures; crowded settings; recapping habits	Safety-engineered devices; no-recapping rule; sharps boxes at arm's reach
Blood & body-fluid exposure	Emergency first assists; unknown patient serostatus; inconsistent glove use	Universal precautions; gloves + eye protection; hepatitis B vaccination
Airborne infection (TB, flu)	Prolonged ward contact; shared rooms; variable respirator fit-testing	Fit-tested respirators; ventilation; prompt isolation protocols
Antineoplastic / chemical agents	Oncology rotations; handling without training	Chemotherapy-competency training; closed-system drug transfer
Radiation	Standing close during fluoroscopy 'to learn', unbadged	Personal dosimeters; distance and shielding; justified standing zones
Sleep loss & fatigue	Night calls plus next-day classes; no duty-hour limits	Enforced rest rules; protected post-call time; scheduling limits
Stress, burnout, bullying	Exams + service demands; power imbalance; stigma around help-seeking	Confidential counselling; anti-bullying policies; mentoring; peer support

Table 2. Principal exposures in clinical training, the student-specific reasons risk is amplified, and the measures shown to reduce them.

6. Prevention: Building a Culture of Safety

The countermeasures for nearly everything described above are known, affordable, and embarrassingly simple. Hepatitis B vaccination — with post-vaccination titre confirmation — converts one of the gravest sharps-injury fears into a manageable risk. Safety-engineered needles and syringes, combined with the iron rule of never recapping, remove the single most common mechanism of injury. A post-exposure protocol that a student can reach within minutes — wash, report, test the source patient, begin prophylaxis where indicated — is the difference between a frightening afternoon

and a life sentence of anxiety. For the mind, the prescriptions are equally concrete: confidential mental-health services that students actually trust, duty-hour limits honoured rather than merely published, and a culture in which seniors model the asking of help.

Yet the deepest prevention is cultural. A teaching hospital that audits student exposures with the same seriousness it audits central-line infections is telling its learners that their safety is part of the curriculum. Institutions that welcome incident reports instead of punishing them, that fit-test respirators for students and not only for staff, that teach safe patient handling as rigorously as they teach anatomy — these are the places where the next generation of physicians learns, from day one, that caring for oneself is not the opposite of caring for patients. It is a prerequisite.




Figure 3. Standard prevention and post-exposure protocol for sharps injuries: engineered devices, PPE, vaccination, immediate first aid, and prompt reporting for post-exposure prophylaxis. (Infographic: Campus Vygon.)

7. Review of the Literature

The modern evidence base on these hazards rests on several sturdy pillars. Global burden estimates for sharps injuries, notably the work of Prüss-Üstün and colleagues published in the Bulletin of the World Health Organization, established that percutaneous exposures among health-care workers number in the millions annually, with the majority occurring in low- and middle-income countries where protective

systems are weakest. Surgical cohort studies have sharpened the picture: Makary and co-workers, following surgeons in training in the United States, documented how frequently trainees are injured and how rarely their injuries are formally reported, even in well-resourced systems.

On the psychological side, the systematic reviews of Dyrbye and colleagues and, later, the meta-analysis by Rotenstein and co-workers in JAMA quantified what students had long said in corridors — that depressive symptoms affect roughly a quarter to a third of medical students worldwide, with burnout, anxiety, and suicidal ideation travelling close behind. Management guidance converges from public-health institutions: the World Health Organization and national public-health services publish exposure-management protocols and infection-prevention standards that apply with equal force to trainees. Read together, the literature tells a consistent and quietly optimistic story — the hazards are well characterized, the countermeasures are known, and the remaining gap is one of implementation and culture rather than knowledge.

8. Conclusion

Medical students enter clinical training to learn how to protect others; the least a teaching hospital can do in return is teach them — systematically, explicitly, and by example — how to protect themselves. The hazards of the clinical years are real and, at times, severe: infected needles, invisible viruses, toxic chemicals, ionizing radiation, wrecked sleep, and a burden of psychological strain that too many students carry in silence. None of this should be romanticized as a rite of passage, because rites of passage are supposed to end. The way forward is neither mysterious nor expensive: vaccination, engineered safety devices, honest reporting, enforced rest, accessible mental-health care, and a hierarchy humble enough to listen to its most junior members. The students who walk through hospital doors tomorrow are the workforce that will staff them for the next forty years. Their safety is not a student issue. It is the future of the profession, measured in unbroken skin, unbroken sleep, and unbroken spirits.

References

1. Prüss-Üstün A, Rapiti E, Hutin Y. Sharps injuries: global burden of disease from sharps injuries to health-care workers. Bull World Health Organ. 2005;83(11):858-63.
2. Makary MA, Al-Attar A, Holzmüller CG, Sexton JB, Syin D, Gilson MM, et al. Needlestick injuries among surgeons in training. N Engl J Med. 2007;356(26):2693-9.

3. Dyrbye LN, Thomas MR, Shanafelt TD. Systematic review of depression, anxiety, and other indicators of psychological distress among U.S. and Canadian medical students. *Acad Med*. 2006;81(4):354-73.
4. Rotenstein LS, Ramos MA, Torre M, Segal JB, Peluso MJ, Guille C, et al. Prevalence of depression, depressive symptoms, and suicidal ideation among medical students: a systematic review and meta-analysis. *JAMA*. 2016;316(21):2214-36.
5. Centers for Disease Control and Prevention. Updated U.S. Public Health Service guidelines for the management of occupational exposures to HIV and recommendations for postexposure prophylaxis. *MMWR Recomm Rep*. 2005;54(RR-9):1-17.
6. World Health Organization. WHO guidelines on drawing blood: best practices in phlebotomy. Geneva: WHO; 2010.
7. World Health Organization. Global health sector strategies on HIV, viral hepatitis and sexually transmitted infections, 2022-2030. Geneva: WHO; 2022.
8. Fisman DN, Harris AD, Rubin MA, Sorock GS, Mittleman MA. Sharps-related injuries in health care workers: a review. *J Occup Environ Med*. 2007;49(10):1089-96.
9. Ishak WW, Lederer S, Mandili C, Nikraves R, Seligman L, Vasa M, et al. Burnout during residency training: a literature review. *J Grad Med Educ*. 2009;1(2):236-42.
10. Wilburn SQ. Needlestick and sharps injury prevention. *Online J Issues Nurs*. 2004;9(3).
11. Shanafelt TD, Balch CM, Bechamps GJ, Russell T, Dyrbye L, Satele D, et al. Burnout and career satisfaction among American surgeons. *Ann Surg*. 2009;250(3):463-71.